

A service of the U.S. National Library of Medicine
and the National Institutes of HealthMy NCBI ?
[Sign In] [Register]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed

for The American journal of cardiology[Jour] AND 762

Go

Clear

Advanced Search

Save Search

Limits

Preview/Index

History

Clipboard

Details

Display

AbstractPlus

Show

20

Sort By

Send to

All: 1

Review: 0

1: Am J Cardiol. 1984 Oct 1;54(7):762-4.

ELSEVIER Links
FULL-TEXT ARTICLE**Prevalence and prognosis of exercise-induced nonsustained ventricular tachycardia in apparently healthy volunteers.****Fleg JL, Lakatta EG.**

Although exercise-induced ventricular tachycardia (VT), whether sustained or nonsustained, is usually associated with significant organic heart disease, its prevalence, associated characteristics and prognostic significance in an asymptomatic, unreferred community-dwelling population are unknown. Therefore, the prevalence of VT associated with maximal treadmill exercise was assessed in 597 male and 325 female volunteers, aged 21 to 96 years (mean +/- standard deviation 54 +/- 16), from the Baltimore Longitudinal Study on Aging who were without apparent heart disease. Ten subjects, 7 men and 3 women, with exercise-induced VT were identified, representing 1.1% of those tested; only 1 was younger than 65 years. All episodes of VT were asymptomatic and nonsustained. In 9 of 10 subjects, VT developed at or near peak exercise. The longest run of VT was 6 beats; multiple runs of VT were present in 4 subjects. Two subjects had exercise-induced ST-segment depression, but subsequent exercise thallium scintigraphic results were negative in each. Compared with a group of age- and sex-matched control subjects, those with asymptomatic, nonsustained VT displayed no difference in exercise duration, maximal heart rate, or the prevalence of coronary risk factors or exercise-induced ischemia as measured by electrocardiography and thallium scintigraphy. Over a mean follow-up period of 2 years, no subject has developed symptoms of heart disease or experienced syncope or sudden death. Thus, exercise-induced VT in apparently healthy subjects occurs almost exclusively in the elderly, is limited to short, asymptomatic runs of 3 to 6 beats usually near peak exercise, and does not portend increased cardiovascular morbidity or mortality rates over a 2-year period of observation.

PMID: 6486025 [PubMed - indexed for MEDLINE]

Related articles

Prevalence and long-term significance of exercise-induced frequent or repetitive ventricular tachycardia. [Am J Cardiol. 1989]

Prevalence and prognostic significance of exercise-induced supraventricular tachycardia in apparently healthy subjects. [Am J Cardiol. 1995]

Independent prognostic significance of ischemic ST-segment response limited to recovery from exercise. [Am J Cardiol. 1998]

Review Evaluation of cardiac arrhythmias by exercise testing. [Herz. 1990]

Review [Tachycardia-induced syncope]. [Ther Umsch. 1997]

» See reviews... | » See all...

Cited by PubMed Central articles

Ventricular tachycardia during exercise testing as a predictor of sudden death in patients with coronary artery disease. [Br Heart J. 1995]

Recent Activity

Turn Off Clear

Prevalence and prognosis of exercise-induced nonsustained ventricular tachycardia in appar...

The American journal of c... (1)

S-T segment elevation during exercise: electrocardiographic and arteriographic

The American journal of... (1)

PubMed

Display

AbstractPlus

Show

20

Sort By

Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer